

RELATIVISTIC KRAMERS–PASTERNAK RECURRENCE RELATIONS

SERGEI K. SUSLOV

ABSTRACT. Recently we have evaluated the matrix elements $\langle Or^p \rangle$, where $O = \{1, \beta, i\alpha\mathbf{n}\beta\}$ are the standard Dirac matrix operators and the angular brackets denote the quantum-mechanical average for the relativistic Coulomb problem, in terms of generalized hypergeometric functions ${}_3F_2(1)$ for all suitable powers and established two sets of Pasternack-type matrix identities for these integrals. The corresponding Kramers–Pasternack type three-term vector recurrence relations are derived here.

1. INTRODUCTION

Recent experimental and theoretical progress has renewed interest in quantum electrodynamics of atomic hydrogenlike systems (see, for example, [22], [23], [27], [28], [36], [53], and [55] and references therein). In the last decade, the two-time Green’s function method of deriving formal expressions for the energy shift of a bound-state level of high- Z few-electron systems was developed [53] and numerical calculations of QED effects in heavy ions were performed with excellent agreement to current experimental data [22], [23], [55]. These advances motivate, among other technical things, investigation of the expectation values of the Dirac matrix operators between the bound-state relativistic Coulomb wave functions. Special cases appear in calculations of the magnetic dipole hyperfine splitting, the electric quadrupole hyperfine splitting, the anomalous Zeeman effect, and the relativistic recoil corrections in hydrogenlike ions (see, for example, [1], [52], [54], [57] and references therein).

In the previous paper [57], the matrix elements $\langle Or^p \rangle$, where $O = \{1, \beta, i\alpha\mathbf{n}\beta\}$ are the standard Dirac matrix operators and the angular brackets denote the quantum-mechanical average for the relativistic Coulomb problem, have been evaluated as sums of three special generalized hypergeometric functions ${}_3F_2(1)$ (or Chebyshev polynomials of a discrete variable) for all suitable powers. As a result, two sets of Pasternack-type matrix symmetry relations for these integrals, when $p \rightarrow -p - 1$ and $p \rightarrow -p - 3$, have been derived. We concentrate on what are essentially radial integrals since, for problems involving spherical symmetry, one can reduce all expectation values to radial integrals by use of the properties of angular momentum.

Our next goal is to establish relativistic analogs of the Kramers–Pasternack recurrence relation [31], [44], and [45] (some progress in this direction had been made in [2]). Here several three-term vector recurrence relations are obtained, which follow in a natural way from the well-known theory of classical orthogonal polynomials of a discrete variable [39] and basic matrix algebra. This paper is organized as follows. We review the nonrelativistic case in the next section and then present

Date: September 30, 2009.

1991 Mathematics Subject Classification. Primary 81Q05. Secondary 33C20.

Key words and phrases. The Dirac equation, hydrogenlike ions, expectation values, Hahn polynomials, Kramers–Pasternack recurrence relation.

relativistic extensions, that seem to be new, in sections 3 and 4. The last section contains a new interpretation of the known two-term recurrence relations for these relativistic expectation values [50], [51]. The new recurrence relations are summarized in appendix A in matrix form for the benefit of the reader.

The author believes that these mathematical results are not only natural and elegant but also will be useful in the current theory of hydrogenlike heavy ions and other exotic relativistic Coulomb systems. At the same time, the mathematical structure behind these expectation values remains not fully understood [2]. For example, to the best of my knowledge, numerous recurrence relations for the radial integrals, obtained with the help of a hypervirial theorem (see [1], [50], [51], [54] and references therein), do not follow in an obvious way from the advanced theory of generalized hypergeometric functions [6]. After more than 80 years of a thorough investigation, the relativistic Coulomb problem keeps generating some mathematical challenges.

2. EXPECTATION VALUES $\langle r^p \rangle$ FOR NONRELATIVISTIC COULOMB PROBLEM

Evaluation of the matrix elements $\langle r^p \rangle$ between nonrelativistic bound-state hydrogenlike wave functions has a long history in quantum mechanics. An incomplete list of references includes [2], [3], [7], [8], [9], [10], [11], [12], [13], [15], [18], [19], [25], [31], [32], [33], [34], [35], [37], [42], [44], [45], [47], [48], [51], [56], [58], [59], [63], and [65]. Different methods were used in order to derive these matrix elements. For example, in Ref. [58] (see also [44], [45]) the mean values for states of definite energy have been obtained in terms of the Chebyshev polynomials of a discrete variable $t_m(x, N) = h_m^{(0,0)}(x, N)$ originally introduced in Refs. [60] and [61]. The so-called Hahn polynomials, introduced also by P. L. Chebyshev [62] and given by

$$h_m^{(\alpha, \beta)}(x, N) = \frac{(1-N)_m (\beta+1)_m}{m!} {}_3F_2 \left(\begin{matrix} -m, \alpha + \beta + m + 1, -x \\ \beta + 1, 1 - N \end{matrix} ; 1 \right), \quad (2.1)$$

were rediscovered and generalized in the late 1940s by W. Hahn (we use the standard definition of the generalized hypergeometric series throughout the paper [6], [21]).

The final results have the following closed forms

$$\langle r^{k-1} \rangle = \frac{1}{2n} \left(\frac{na_0}{2Z} \right)^{k-1} t_k(n-l-1, -2l-1), \quad (2.2)$$

when $k = 0, 1, 2, \dots$ and

$$\left\langle \frac{1}{r^{k+2}} \right\rangle = \frac{1}{2n} \left(\frac{2Z}{na_0} \right)^{k+2} t_k(n-l-1, -2l-1), \quad (2.3)$$

when $k = 0, 1, \dots, 2l$. Here $a_0 = \hbar^2/me^2$ is the Bohr radius (more details can be found in Refs. [14] and [58]).

The ease of handling of these matrix elements for the discrete levels is greatly increased if use is made of the known properties of these classical polynomials of Chebyshev [21], [26], [29], [39], [40] and [60], [61], [62]. The direct consequences of (2.2)–(2.3) are an inversion relation:

$$\left\langle \frac{1}{r^{k+2}} \right\rangle = \left(\frac{2Z}{na_0} \right)^{2k+1} \frac{(2l-k)!}{(2l+k+1)!} \langle r^{k-1} \rangle \quad (2.4)$$

with $0 \leq k \leq 2l$ and the three-term recurrence relation:

$$\langle r^k \rangle = \frac{2n(2k+1)}{k+1} \left(\frac{na_0}{2Z} \right) \langle r^{k-1} \rangle - \frac{k((2l+1)^2 - k^2)}{k+1} \left(\frac{na_0}{2Z} \right)^2 \langle r^{k-2} \rangle \quad (2.5)$$

with the initial conditions

$$\left\langle \frac{1}{r} \right\rangle = \frac{Z}{a_0 n^2}, \quad \langle 1 \rangle = 1, \quad (2.6)$$

which is convenient for evaluation of the mean values $\langle r^k \rangle$ for $k \geq 1$.

In our approach, the recurrence relation (2.5) is a special case of the three-term recurrence relation for the Hahn polynomials (3.8). It was originally found by Kramers and Pasternack in the late 1930s [31], [44], and [45]. The inversion symmetry (2.4), which is also due to Pasternack, has been rediscovered many years later [12], [37] (see also [24] and [25] for historical comments). Generalizations of (2.4)–(2.5) for off-diagonal matrix elements are discussed in Refs. [11], [17], [20], [25], [38], [41], [42], [46], [49], [50], [51], [56], and [59]. The properties of the hydrogenlike radial matrix elements are considered from a group-theoretical viewpoint in Refs. [3], [13], and [42]. Extensions to the relativistic case are given in [1], [2], [16], [19], [43], [57], and [58] (see also references therein and the following sections of this paper).

In a retrospect, Pasternack's papers [44], [45] had paved the road to the discovery of the continuous Hahn polynomials in the mid 1980s (see [4], [5], [30], [58] and references therein).

3. THREE-TERM RECURRENCE RELATIONS FOR RELATIVISTIC MATRIX ELEMENTS

In the present paper we establish relativistic analogs of the Kramers–Pasternack recurrence relations (2.5) for the following set of integrals of the relativistic radial functions:

$$A_p = \int_0^\infty r^{p+2} (F^2(r) + G^2(r)) dr, \quad (3.1)$$

$$B_p = \int_0^\infty r^{p+2} (F^2(r) - G^2(r)) dr, \quad (3.2)$$

$$C_p = \int_0^\infty r^{p+2} F(r) G(r) dr. \quad (3.3)$$

With the notations from Refs. [57] and [58], one can evaluate these integrals in terms of the Chebyshev and Hahn polynomials of a discrete variable and present the answer, say, when $p \geq 0$, in the following closed form:

$$4\mu\nu^2 (2a\beta)^p A_p \quad (3.4)$$

$$\begin{aligned} &= a\kappa (\varepsilon\kappa + \nu) h_{p+1}^{(0,0)}(n-1, -1-2\nu) \\ &- 2\frac{p+2}{p+1} \mu (\varepsilon^2\kappa^2 - \nu^2) h_p^{(1,1)}(n-1, -1-2\nu) \\ &+ a\kappa (\varepsilon\kappa - \nu) h_{p+1}^{(0,0)}(n, 1-2\nu), \end{aligned}$$

$$4\mu\nu (2a\beta)^p B_p \quad (3.5)$$

$$= a (\varepsilon\kappa + \nu) h_{p+1}^{(0,0)}(n-1, -1-2\nu)$$

$$\begin{aligned}
& -a(\varepsilon\kappa - \nu) h_{p+1}^{(0,0)}(n, 1 - 2\nu), \\
& 8\mu\nu^2 (2a\beta)^p C_p \\
& = a\mu(\varepsilon\kappa + \nu) h_{p+1}^{(0,0)}(n - 1, -1 - 2\nu) \\
& - 2\frac{p+2}{p+1}\kappa(\varepsilon^2\kappa^2 - \nu^2) h_p^{(1,1)}(n - 1, -1 - 2\nu) \\
& + a\mu(\varepsilon\kappa - \nu) h_{p+1}^{(0,0)}(n, 1 - 2\nu).
\end{aligned} \tag{3.6}$$

Here,

$$\begin{aligned}
\kappa &= \pm(j + 1/2), & \nu &= \sqrt{\kappa^2 - \mu^2}, \\
\mu &= \alpha Z = Ze^2/\hbar c, & a &= \sqrt{1 - \varepsilon^2}, \\
\varepsilon &= E/mc^2, & \beta &= mc/\hbar
\end{aligned} \tag{3.7}$$

with the total angular momentum $j = 1/2, 3/2, 5/2, \dots$ (see [57] and [58] for more details).

Although a set of useful recurrence relations between the relativistic matrix elements was derived by Shabaev [50], [51] (see also [1], [19], [54], [57], [64], and the last section of this paper) on the basis of a hypervirial theorem, the corresponding relativistic Kramers–Pasternack type relations seem to be missing in the available literature. Our equations (3.4)–(3.6) reveal that they should have a vector form.

Here, one can apply a familiar three-term recurrence relation for the Hahn polynomials [39], [40],

$$xh_m^{(\alpha, \beta)}(x, N) = \alpha_m h_{m+1}^{(\alpha, \beta)}(x, N) + \beta_m h_m^{(\alpha, \beta)}(x, N) + \gamma_m h_{m-1}^{(\alpha, \beta)}(x, N) \tag{3.8}$$

with

$$\alpha_m = \frac{(m+1)(\alpha + \beta + m + 1)}{(\alpha + \beta + 2m + 1)(\alpha + \beta + 2m + 2)}, \tag{3.9}$$

$$\beta_m = \frac{\alpha - \beta + 2N - 2}{4} + \frac{(\beta^2 - \alpha^2)(\alpha + \beta + 2N)}{4(\alpha + \beta + 2m)(\alpha + \beta + 2m + 2)}, \tag{3.10}$$

$$\gamma_m = \frac{(\alpha + m)(\beta + m)(\alpha + \beta + N + m)(N - m)}{(\alpha + \beta + 2m)(\alpha + \beta + 2m + 1)} \tag{3.11}$$

three times for the corresponding special cases in (3.4)–(3.6). Introducing two sets of vectors

$$\mathbf{A}_p = \begin{pmatrix} A_p \\ B_p \\ C_p \end{pmatrix}, \quad \mathbf{X}_p = \begin{pmatrix} h_{p+1}^{(0,0)}(n - 1, -1 - 2\nu) \\ \frac{p+2}{p+1}h_p^{(1,1)}(n - 1, -1 - 2\nu) \\ h_{p+1}^{(0,0)}(n, 1 - 2\nu) \end{pmatrix}, \tag{3.12}$$

we conclude that the recurrence relations in question should have the following matrix structure:

$$\mathbf{A}_p = T\mathbf{X}_p, \quad \mathbf{X}_p = T^{-1}\mathbf{A}_p \tag{3.13}$$

with

$$\mathbf{X}_{p+1} = U_p\mathbf{X}_p + V_p\mathbf{X}_{p-1}, \tag{3.14}$$

where U_p and W_p are known diagonal matrices. Then

$$\mathbf{A}_{p+1} = D_p\mathbf{A}_p + E_p\mathbf{A}_{p-1}, \tag{3.15}$$

where

$$D_p = TU_p T^{-1}, \quad E_p = TV_p T^{-1} \quad (3.16)$$

are similar matrices.

The elementary linear algebra argument outlined above allows us to obtain a relativistic generalization of the Kramers–Pasternack recurrence relation (2.5) as follows:

$$\begin{aligned} (2a\beta)^2 A_{p+1} &= 4\beta\varepsilon\mu \frac{2p+3}{p+2} A_p \\ &\quad - \frac{p(p+2)(4\nu^2 - (p+1)^2) + 4\kappa^2}{(p+1)(p+2)} A_{p-1} \\ &\quad - 4\kappa \frac{p+1}{p+2} B_{p-1} + \frac{8\kappa\mu}{(p+1)(p+2)} C_{p-1}, \end{aligned} \quad (3.17)$$

$$\begin{aligned} (2a\beta)^2 B_{p+1} &= 4\beta\varepsilon\mu \frac{2p+3}{p+2} B_p \\ &\quad - (4\nu^2 - p(p+2)) \frac{p+1}{p+2} B_{p-1} \\ &\quad - 4\kappa \frac{p+1}{p+2} A_{p-1} + 8\mu \frac{p+1}{p+2} C_{p-1}, \end{aligned} \quad (3.18)$$

$$\begin{aligned} (2a\beta)^2 C_{p+1} &= 4\beta\varepsilon\mu \frac{2p+3}{p+2} C_p \\ &\quad - \frac{2\kappa\mu}{(p+1)(p+2)} A_{p-1} - 2\mu \frac{p+1}{p+2} B_{p-1} \\ &\quad - \frac{p(p+2)(4\nu^2 - (p+1)^2) - 4\mu^2}{(p+1)(p+2)} C_{p-1}. \end{aligned} \quad (3.19)$$

Our relations (3.17)–(3.19), together with the symmetry under the reflections, $p \rightarrow -p-1$ and $p \rightarrow -p-3$, established in [57], give a possibility to compute all the required matrix elements in a recursive manner without direct evaluation of the integrals. The corresponding initial values (A.3)–(A.4) can be found in Ref. [57] or elsewhere.

A different representation of these integrals in terms of Chebyshev polynomials is also available [57]:

$$\begin{aligned} 2a\mu (2a\beta)^p A_p &= a(\mu + a\kappa) h_p^{(0,0)}(n-1, -2\nu) \\ &\quad + 2\varepsilon(\mu^2 - a^2\kappa^2) h_{p-1}^{(1,1)}(n-1, -1-2\nu) \\ &\quad + a(\mu - a\kappa) h_p^{(0,0)}(n, -2\nu), \end{aligned} \quad (3.20)$$

$$\begin{aligned} 2a\mu (2a\beta)^p B_p &= a\varepsilon(\mu + a\kappa) h_p^{(0,0)}(n-1, -2\nu) \\ &\quad + 2(\mu^2 - a^2\kappa^2) h_{p-1}^{(1,1)}(n-1, -1-2\nu) \end{aligned} \quad (3.21)$$

$$\begin{aligned}
& +a\varepsilon (\mu - a\kappa) \, h_p^{(0,0)}(n, -2\nu), \\
& 4\mu (2a\beta)^p C_p \\
& = a\varepsilon (\mu + a\kappa) \, h_p^{(0,0)}(n-1, -2\nu) \\
& - a\varepsilon (\mu - a\kappa) \, h_p^{(0,0)}(n, -2\nu),
\end{aligned} \tag{3.22}$$

when $p \geq 1$. A similar matrix manipulation results in another recurrence relation:

$$\begin{aligned}
(2a\beta)^2 A_{p+1} &= 4\beta\varepsilon\mu \frac{(a^2(p+1)^2 - 1)(2p+1)}{a^2p(p+1)(p+2)} A_p \\
&+ \frac{4\beta\varepsilon^2\mu(2p+1)}{a^2p(p+1)(p+2)} B_p - 4\beta\frac{2p+1}{p+1} C_p \\
&- \frac{(a^2(p+1)^2 - 1)(4\nu^2 - p^2)}{a^2(p+1)(p+2)} A_{p-1} - \frac{\varepsilon(4\nu^2 - p^2)}{a^2(p+1)(p+2)} B_{p-1},
\end{aligned} \tag{3.23}$$

$$\begin{aligned}
(2a\beta)^2 B_{p+1} &= -\frac{4\beta\varepsilon^2\mu(2p+1)}{a^2p(p+1)(p+2)} A_p \\
&+ 4\beta\varepsilon\mu \frac{(a^2(p+1)^2 + \varepsilon^2)(2p+1)}{a^2p(p+1)(p+2)} B_p - 4\beta\varepsilon\frac{2p+1}{p+1} C_p \\
&+ \frac{\varepsilon(4\nu^2 - p^2)}{a^2(p+1)(p+2)} A_{p-1} - \frac{(4\nu^2 - p^2)(a^2(p+1)^2 + \varepsilon^2)}{a^2(p+1)(p+2)} B_{p-1},
\end{aligned} \tag{3.24}$$

$$\begin{aligned}
(2a\beta)^2 C_{p+1} &= -\beta\frac{2p+1}{p+1} A_p + \beta\varepsilon\frac{2p+1}{p+1} B_p \\
&+ 4\beta\varepsilon\mu\frac{2p+1}{p+1} C_p - (4\nu^2 - p^2)\frac{p}{p+1} C_{p-1}.
\end{aligned} \tag{3.25}$$

These recurrence relations are summarized for the benefit of the reader in matrix form in appendix A. One should take special values (A.4)–(A.5) as the initial data. A direct derivation of the relativistic three-term vector recurrence relations (3.17)–(3.19) and (3.23)–(3.25) on the basis of a hypervirial theorem needs to be found.

4. INDEPENDENT INTEGRALS

The integrals (3.1)–(3.3) are linearly dependent:

$$(2\kappa + \varepsilon(p+1)) A_p - (2\varepsilon\kappa + p+1) B_p = 4\mu C_p \tag{4.1}$$

(see, for example, [1], [50], [51], and [57] for more details). Thus, eliminating C_p , say from (3.23)–(3.24), we obtain the following three-term vector recurrence relation between the integrals A_p and B_p only:

$$\begin{aligned}
(2a\beta)^2 A_{p+1} & \\
&= (2p+1)\beta \frac{4\varepsilon\mu^2(a^2(p+1)^2 - 1) - a^2p(p+2)(2\kappa + \varepsilon(p+1))}{a^2\mu p(p+1)(p+2)} A_p
\end{aligned} \tag{4.2}$$

$$\begin{aligned}
& + (2p+1) \beta \frac{4\varepsilon^2 \mu^2 + a^2 p(p+2)(2\varepsilon\kappa + p+1)}{a^2 \mu p(p+1)(p+2)} B_p \\
& - \frac{(a^2(p+1)^2 - 1)(4\nu^2 - p^2)}{a^2(p+1)(p+2)} A_{p-1} - \frac{\varepsilon(4\nu^2 - p^2)}{a^2(p+1)(p+2)} B_{p-1}
\end{aligned}$$

and

$$\begin{aligned}
(2a\beta)^2 B_{p+1} & = - (2p+1) \beta \varepsilon \frac{4\varepsilon \mu^2 + a^2 p(p+2)(2\kappa + \varepsilon(p+1))}{a^2 \mu p(p+1)(p+2)} A_p \\
& + (2p+1) \beta \varepsilon \frac{4\mu^2(a^2(p+1)^2 + \varepsilon^2) + a^2 p(p+2)(2\varepsilon\kappa + p+1)}{a^2 \mu p(p+1)(p+2)} B_p \\
& + \frac{\varepsilon(4\nu^2 - p^2)}{a^2(p+1)(p+2)} A_{p-1} - \frac{(4\nu^2 - p^2)(a^2(p+1)^2 + \varepsilon^2)}{a^2(p+1)(p+2)} B_{p-1}.
\end{aligned} \tag{4.3}$$

In a similar fashion, from (3.17)–(3.18):

$$\begin{aligned}
(2a\beta)^2 A_{p+1} & = 4\beta\varepsilon\mu \frac{2p+3}{p+2} A_p \\
& + p \frac{2\varepsilon\kappa - (p+2)(4\nu^2 - (p+1)^2)}{(p+1)(p+2)} A_{p-1} \\
& - 2\kappa \frac{2\varepsilon\kappa - 1 + (p+1)(2p+3)}{(p+1)(p+2)} B_{p-1}
\end{aligned} \tag{4.4}$$

and

$$\begin{aligned}
(2a\beta)^2 B_{p+1} & = 4\beta\varepsilon\mu \frac{2p+3}{p+2} B_p \\
& + 2\varepsilon p \frac{p+1}{p+2} A_{p-1} - (4(\varepsilon\kappa + \nu^2) - p^2) \frac{p+1}{p+2} B_{p-1}.
\end{aligned} \tag{4.5}$$

Matrix forms of these identities and the corresponding initial values are given in appendix A.

5. TWO-TERM RECURRENCE RELATIONS

The three-term recurrence relations for the relativistic radial integrals, examples of which have been found in the previous sections, are obviously not unique. Moreover, if

$$\begin{aligned}
\mathbf{A}_{p+1} & = D_p^{(1)} \mathbf{A}_p + E_p^{(1)} \mathbf{A}_{p-1}, \\
\mathbf{A}_{p+1} & = D_p^{(2)} \mathbf{A}_p + E_p^{(2)} \mathbf{A}_{p-1},
\end{aligned} \tag{5.1}$$

then

$$\begin{aligned}
(\alpha_p + \beta_p) \mathbf{A}_{p+1} & = (\alpha_p D_p^{(1)} + \beta_p D_p^{(2)}) \mathbf{A}_p \\
& + (\alpha_p E_p^{(1)} + \beta_p E_p^{(2)}) \mathbf{A}_{p-1}
\end{aligned} \tag{5.2}$$

for two arbitrary sequences of scalars α_p and β_p . One special case is of a particular interest.

Subtracting (4.2)–(4.3) from (4.4)–(4.5) one gets the following matrix equation

$$P_p \begin{pmatrix} A_p \\ B_p \end{pmatrix} = Q_p \begin{pmatrix} A_{p-1} \\ B_{p-1} \end{pmatrix} \quad (5.3)$$

with

$$\det P_p = -8\beta^2\varepsilon \frac{\kappa a^2 (p+2)(2p+1) + 2\varepsilon\mu^2}{p(p+1)(p+2)^2}, \quad (5.4)$$

$$\det Q_p = -2\varepsilon(4\nu^2 - p^2) \frac{\kappa a^2 (p+2)(2p+1) + 2\varepsilon\mu^2}{a^2(p+2)^2(p+1)^2} \quad (5.5)$$

by a computer algebra system. We have omitted the explicit forms of the P and Q matrices and their inverses. It should be easily done by the reader.

The two-term recurrence solutions of the form

$$\begin{pmatrix} A_p \\ B_p \end{pmatrix} = S_p \begin{pmatrix} A_{p-1} \\ B_{p-1} \end{pmatrix}, \quad \begin{pmatrix} A_{p-1} \\ B_{p-1} \end{pmatrix} = S_p^{-1} \begin{pmatrix} A_p \\ B_p \end{pmatrix} \quad (5.6)$$

where found by Shabaev [50], [51] by a different method. In our notations,

$$\begin{aligned} A_{p+1} = & -(p+1) \frac{4\nu^2\varepsilon + 2\kappa(p+2) + \varepsilon(p+1)(2\kappa\varepsilon + p+2)}{4(1-\varepsilon^2)(p+2)\beta\mu} A_p \\ & + \frac{4\mu^2(p+2) + (p+1)(2\kappa\varepsilon + p+1)(2\kappa\varepsilon + p+2)}{4(1-\varepsilon^2)(p+2)\beta\mu} B_p, \end{aligned} \quad (5.7)$$

$$\begin{aligned} B_{p+1} = & -(p+1) \frac{4\nu^2 + 2\kappa\varepsilon(2p+3) + \varepsilon^2(p+1)(p+2)}{4(1-\varepsilon^2)(p+2)\beta\mu} A_p \\ & + \frac{4\mu^2\varepsilon(p+2) + (p+1)(2\kappa\varepsilon + p+1)(2\kappa + \varepsilon(p+2))}{4(1-\varepsilon^2)(p+2)\beta\mu} B_p \end{aligned} \quad (5.8)$$

and

$$\begin{aligned} A_{p-1} = & \beta \frac{4\mu^2\varepsilon(p+1) + p(2\kappa\varepsilon + p)(2\kappa + \varepsilon(p+1))}{\mu(4\nu^2 - p^2)p} A_p \\ & - \beta \frac{4\mu^2(p+1) + p(2\kappa\varepsilon + p)(2\kappa\varepsilon + p+1)}{\mu(4\nu^2 - p^2)p} B_p, \end{aligned} \quad (5.9)$$

$$\begin{aligned} B_{p-1} = & \beta \frac{4\nu^2 + 2\kappa\varepsilon(2p+1) + \varepsilon^2p(p+1)}{\mu(4\nu^2 - p^2)} A_p \\ & - \beta \frac{4\nu^2\varepsilon + 2\kappa(p+1) + \varepsilon p(2\kappa\varepsilon + p+1)}{\mu(4\nu^2 - p^2)} B_p, \end{aligned} \quad (5.10)$$

respectively.

We have shown that these solutions can be derived from the three-term vector recurrence relations found in this paper. As a by-product, the factorization of Shabaev's matrices, namely,

$$S_p = P_p^{-1} Q_p, \quad S_p^{-1} = Q_p^{-1} P_p, \quad (5.11)$$

is given. This can be directly verified with the help of a computer algebra system. Then

$$\det S_p = \frac{\det Q_p}{\det P_p} = \frac{(4\nu^2 - p^2)p}{4(a\beta)^2(p+1)}. \quad (5.12)$$

Further details are left to the reader.

Acknowledgments. I thank Carlos Castillo-Chávez for support and encouragement and David Murillo for help. I am grateful to Vladimir M. Shabaev for a copy of [50]. The referees' suggestions are also very much appreciated.

APPENDIX A. MATRIX FORM OF THREE-TERM RECURRENCE RELATIONS

The matrix structures of our recurrence relations (3.17)–(3.19) and (3.23)–(3.25) are given by

$$(2a\beta)^2 \begin{pmatrix} A_{p+1} \\ B_{p+1} \\ C_{p+1} \end{pmatrix} = 4\beta\varepsilon\mu \frac{2p+3}{p+2} \begin{pmatrix} A_p \\ B_p \\ C_p \end{pmatrix} \quad (A.1)$$

$$- \begin{pmatrix} \frac{p(p+2)(4\nu^2 - (p+1)^2) + 4\kappa^2}{(p+1)(p+2)} & 4\kappa \frac{p+1}{p+2} & -\frac{8\kappa\mu}{(p+1)(p+2)} \\ 4\kappa \frac{p+1}{p+2} & (4\nu^2 - p(p+2)) \frac{p+1}{p+2} & -8\mu \frac{p+1}{p+2} \\ \frac{2\kappa\mu}{(p+1)(p+2)} & 2\mu \frac{p+1}{p+2} & \frac{p(p+2)(4\nu^2 - (p+1)^2) - 4\mu^2}{(p+1)(p+2)} \end{pmatrix} \begin{pmatrix} A_{p-1} \\ B_{p-1} \\ C_{p-1} \end{pmatrix}$$

and

$$(2a\beta)^2 \begin{pmatrix} A_{p+1} \\ B_{p+1} \\ C_{p+1} \end{pmatrix} \quad (A.2)$$

$$= \beta \frac{2p+1}{p+1} \begin{pmatrix} \frac{4\varepsilon\mu(a^2(p+1)^2 - 1)}{a^2 p(p+2)} & \frac{4\varepsilon^2\mu}{a^2 p(p+2)} & -4 \\ -\frac{4\varepsilon^2\mu}{a^2 p(p+2)} & \frac{4\varepsilon\mu(a^2(p+1)^2 + \varepsilon^2)}{a^2 p(p+2)} & -4\varepsilon \\ -1 & \varepsilon & 4\varepsilon\mu \end{pmatrix} \begin{pmatrix} A_p \\ B_p \\ C_p \end{pmatrix}$$

$$- \frac{(4\nu^2 - p^2)}{p+1} \begin{pmatrix} \frac{(a^2(p+1)^2 - 1)}{a^2(p+2)} & \frac{\varepsilon}{a^2(p+2)} & 0 \\ -\frac{\varepsilon}{a^2(p+2)} & \frac{(a^2(p+1)^2 + \varepsilon^2)}{a^2(p+2)} & 0 \\ 0 & 0 & p \end{pmatrix} \begin{pmatrix} A_{p-1} \\ B_{p-1} \\ C_{p-1} \end{pmatrix},$$

respectively. The initial vectors are

$$\begin{pmatrix} A_{-1} \\ B_{-1} \\ C_{-1} \end{pmatrix} = \begin{pmatrix} \frac{\beta}{\mu\nu} (1 - \varepsilon^2) (\varepsilon\nu + \mu\sqrt{1 - \varepsilon^2}) \\ \frac{\beta a^2}{\mu} \\ \frac{\kappa}{2\mu\nu} a^3 \beta \end{pmatrix}, \quad (A.3)$$

$$\begin{pmatrix} A_0 \\ B_0 \\ C_0 \end{pmatrix} = \begin{pmatrix} 1 \\ \varepsilon \\ \frac{\kappa}{2\mu} (1 - \varepsilon^2) \end{pmatrix}, \quad (A.4)$$

$$\begin{pmatrix} A_1 \\ B_1 \\ C_1 \end{pmatrix} = \begin{pmatrix} \frac{3\varepsilon\mu^2 - \kappa(1-\varepsilon^2)(1+\varepsilon\kappa)}{2\beta\mu(1-\varepsilon^2)} \\ \frac{3\varepsilon^2\mu^2 - (1-\varepsilon^2)(\varepsilon\kappa + \nu^2)}{2\beta\mu(1-\varepsilon^2)} \\ \frac{2\varepsilon\kappa - 1}{4\beta} \end{pmatrix}. \quad (\text{A.5})$$

One should use the initial data (A.3)–(A.4) in the case of the recurrence relation (A.1) and (A.4)–(A.5) for (A.2).

Our relations (4.2)–(4.3) and (4.4)–(4.5) take the following matrix forms:

$$\begin{aligned} (2a\beta)^2 \begin{pmatrix} A_{p+1} \\ B_{p+1} \end{pmatrix} \\ = \beta \frac{2p+1}{p+1} \begin{pmatrix} \frac{4\varepsilon\mu^2(a^2(p+1)^2-1) - a^2p(p+2)(2\kappa+\varepsilon(p+1))}{a^2\mu p(p+2)} & \frac{4\varepsilon^2\mu^2 + a^2p(p+2)(2\varepsilon\kappa+p+1)}{a^2\mu p(p+2)} \\ -\varepsilon \frac{4\varepsilon\mu^2 + a^2p(p+2)(2\kappa+\varepsilon(p+1))}{a^2\mu p(p+2)} & \varepsilon \frac{4\mu^2(a^2(p+1)^2+\varepsilon^2) + a^2p(p+2)(2\varepsilon\kappa+p+1)}{a^2\mu p(p+2)} \end{pmatrix} \\ \times \begin{pmatrix} A_p \\ B_p \end{pmatrix} - \frac{(4\nu^2 - p^2)}{a^2(p+1)(p+2)} \begin{pmatrix} a^2(p+1)^2 - 1 & \varepsilon \\ -\varepsilon & a^2(p+1)^2 + \varepsilon^2 \end{pmatrix} \begin{pmatrix} A_{p-1} \\ B_{p-1} \end{pmatrix} \end{aligned} \quad (\text{A.6})$$

and

$$\begin{aligned} (2a\beta)^2 \begin{pmatrix} A_{p+1} \\ B_{p+1} \end{pmatrix} &= 4\beta\varepsilon\mu \frac{2p+3}{p+2} \begin{pmatrix} A_p \\ B_p \end{pmatrix} \\ &- \begin{pmatrix} -p \frac{2\varepsilon\kappa - (p+2)(4\nu^2 - (p+1)^2)}{(p+1)(p+2)} & 2\kappa \frac{2\varepsilon\kappa - 1 + (p+1)(2p+3)}{(p+1)(p+2)} \\ -2\varepsilon p \frac{p+1}{p+2} & (4(\varepsilon\kappa + \nu^2) - p^2) \frac{p+1}{p+2} \end{pmatrix} \begin{pmatrix} A_{p-1} \\ B_{p-1} \end{pmatrix}, \end{aligned} \quad (\text{A.7})$$

respectively.

REFERENCES

- [1] G. S. Adkins, *Dirac–Coulomb energy levels and expectation values*, Am. J. Phys. **76** (2008) #6, 579–584.
- [2] D. Andrae, *Recursive evaluation of expectation values $\langle r^k \rangle$ for arbitrary states of the relativistic one-electron atom*, J. Phys. B: Atom. Molec. Phys. **30** (1997) #20, 4435–4451.
- [3] L. Armstrong, Jr., *Mean values of powers of the radius for hydrogenic electron orbits*, Phys. Rev. A **3** (1971) #5, 1546–1550.
- [4] R. A. Askey, *Continuous Hahn polynomials*, J. Phys. A: Math. Gen. **18** (1985), L1017–L1019.
- [5] N. M. Atakishyev and S. K. Suslov, *The Hahn and Meixner polynomials of imaginary argument and some of their applications*, J. Phys. A: Math. Gen. **18** (1985), 1583–1596.
- [6] W. N. Bailey, *Generalized Hypergeometric Series*, Cambridge University Press, Cambridge, 1935.
- [7] S. Balasubramanian, *A simple derivation of the recurrence relation for $\langle r^N \rangle$* , Am. J. Phys. **68** (2000) #10, 959–960.
- [8] A. I. Baz, Ya. B. Zel’dovich, and A. M. Perelomov, *Scattering, Reactions and Decays in Nonrelativistic Quantum Mechanics* [in Russian], Nauka, Moscow, 1971.
- [9] H. Beker, *A simple calculation of $\langle 1/r^2 \rangle$ for the hydrogen atom and the three-dimensional harmonic oscillator*, Am. J. Phys. **65** (1997) #11, 1118–1119.
- [10] H. A. Bethe and E. E. Salpeter, *Quantum Mechanics of One- and Two-Electron Atoms*, Springer–Verlag, Berlin, 1957.
- [11] P. Blanchard, *A new recurrence relation for hydrogenic radial matrix elements*, J. Phys. B: At. Mol. Phys. **7** (1974) #9, 993–1005.

- [12] K. Bockasten, *Mean values of powers of the radius for hydrogenic electron orbits*, Phys. Rev. A **9** (1974), 1087–1089.
- [13] E. Chacón, D. Levi, and M. Moshinsky, *Lie algebras in the Schrödinger picture and radial matrix elements*, J. Math. Phys. **17** (1976) #10, 1919–1929.
- [14] R. Cordero-Soto and S. K. Suslov, *Expectation values $\langle r^p \rangle$ for harmonic oscillator in R^n* , arXiv: 0908.0032v3 [math-ph] 6 Aug 2009.
- [15] L. J. Curtis, *Classical mnemonic approach for obtaining hydrogenic expectation values r^p* , Phys. Rev. A **43** (1991) #1, 568–569.
- [16] L. Davis, *A note on the wave functions of the relativistic hydrogen atom*, Phys. Rev. **56** (1939), 186–187.
- [17] Yi-B. Ding, *On the Schrödinger radial ladder operator*, J. Phys. A: Math. Gen. **20** (1987) #18, 6293–6301.
- [18] G. W. F. Drake and R. A. Swainson, *Expectation values of r^p for arbitrary hydrogenic states*, Phys. Rev. A **42** (1990) #3, 1123–1126.
- [19] J. H. Epstein, S. T. Epstein, *Some applications of hypervirial theorems to the calculation of average values*, Am. J. Phys. **30** (1962) #4, 266–268.
- [20] S. T. Epstein, J. H. Epstein, and B. Kennedy, *Recursion relations for Coulomb matrix elements*, J. Math. Phys. **8** (1967) #9, 1747.
- [21] A. Erdélyi, *Higher Transcendental Functions*, Vols. I–III, A. Erdélyi, ed., McGraw–Hill, 1953.
- [22] A. Gumberidze *et al.*, *Quantum electrodynamics in strong electric fields: The ground state Lamb shift in hydrogenlike uranium*, Phys. Rev. Lett. **94** (2005), 223001: 1–4.
- [23] A. Gumberidze *et al.*, *Precision tests of QED in strong fields: experiments on hydrogen- and helium-like uranium*, J. Phys. Conf. Ser. **58** (2007), 87–92.
- [24] J. D. Hey, *On the momentum representation of hydrogenic wave functions: Some properties and an application*, Am. J. Phys. **61** (1993) #1, 28–35.
- [25] J. D. Hey, *Further properties of hydrogenic wave functions*, Am. J. Phys. **61** (1993) #8, 741–749.
- [26] S. Karlin and J. L. McGregor, *The Hahn polynomials, formulas and an application*, Scripta Math. **26** (1961), 33–46.
- [27] S. G. Karshenboim *et al* (Eds.), *The Hydrogen Atom: Precision Physics of Simple Atomic Systems*, Lecture Notes in Physics, Vol. 570, Springer–Verlag, Berlin, Heidelberg, 2001.
- [28] S. G. Karshenboim and V. B. Smirnov (Eds.), *Precision Physics of Simple Atomic Systems*, Lecture Notes in Physics, Vol. 627, Springer–Verlag, Berlin, Heidelberg, 2003.
- [29] R. Koekoek and R. F. Swarttouw, *The Askey scheme of hypergeometric orthogonal polynomials and its q -analogues*, Report 94–05, Delft University of Technology, 1994.
- [30] H. T. Koelink, *On Jacobi and continuous Hahn polynomials*, Proc. Amer. Math. Soc. **124** (1996), 887–898.
- [31] H. A. Kramers, *Die Grundlagen der Quantentheorie/Quantentheorie des Electrons und der Strahlung*, Akademische Verlagsgesellschaft, Leipzig, 1938; Engl. transl., *Quantum Mechanics*, North-Holland, Amsterdam, 1957, section 59, equation (6.74).
- [32] L. D. Landau and E. M. Lifshitz, *Quantum Mechanics: Nonrelativistic Theory*, Pergamon Press, Oxford, 1977.
- [33] R. L. Liboff, *Introduction to Quantum Mechanics*, third edition, Addison–Wesley, Longman, Reading, MA, 1998, pp. 475–476.
- [34] H. Marxer, *Exact corresponding relationship for the expectation values r^{-k} for hydrogenlike states*, Phys. Rev. A **44** (1991) #3, 1543–1548.
- [35] A. Messiah, *Quantum Mechanics*, Vol. 1, North-Holland, Amsterdam, 1961, p. 431.
- [36] P. J. Mohr, G. Plunien, and G. Soff, *QED corrections in heavy atoms*, Phys. Rep. **293** (1998), 227–369.
- [37] R. M. More, *Inversion symmetry of hydrogenic expectation values*, J. Phys. B: At. Mol. Phys. **16** (1993) #21, 3859–3861.
- [38] B. Moreno, A. López Piñeiro, and R. H. Tipping, *Algebraic solutions for the hydrogenic radial Schrödinger equation: matrix elements for arbitrary powers of several r -dependent operators*, J. Phys. A: Math. Gen. **24** (1991) #2, 385–390.
- [39] A. F. Nikiforov, S. K. Suslov, and V. B. Uvarov, *Classical Orthogonal Polynomials of a Discrete Variable*, Springer–Verlag, Berlin, New York, 1991.
- [40] A. F. Nikiforov and V. B. Uvarov, *Special Functions of Mathematical Physics*, Birkhäuser, Basel, Boston, 1988.
- [41] H. N. Núñez-Yépez, J. L. López-Bonilla, and A. L. Salas-Brito, *Generalized hypervirial and recurrence relations for hydrogenic matrix elements*, J. Phys. B: At. Mol. Phys. **28** (1995), L525–L529.

- [42] P. C. Ojha and D. S. F. Crothers, *On a simple relation between hydrogenic radial matrix elements*, J. Phys. B: At. Mol. Phys. **17** (1984) #24, 4797–4800.
- [43] L. C. Owono Owono, M. G. Kwato Njock, and B. Oumarou, *Expectation values of r^q between Dirac and quasirelativistic wave functions in the quantum-defect approximation*, Phys. Rev. A **66** (2002), 052503.
- [44] S. Pasternack, *On the mean value of r^s for Keplerian systems*, Proc. Nat. Acad. Sci. **23** (1937) #2, 91–94; #4, 250 (erratum).
- [45] S. Pasternack, *A generalization of the polynomials $F_n(x)$* , London, Edinburgh, Dublin Philosophical Magazine and J. Science, Ser. 7 **28** (1939), 209–226.
- [46] S. Pasternack and R. M. Sternheimer, *An orthogonality property of hydrogenlike radial functions*, J. Math. Phys. **3** (1962) #6, 1280.
- [47] A. M. Perelomov and Ya. B. Zel'dovich, *Quantum Mechanics: Selected Topics*, World Scientific Publishing Co., Inc., River Edge, NJ, 1998.
- [48] W.-C. Qiang and S.-H. Dong, *An alternative approach to calculation the mean values $\overline{r^k}$ for hydrogen-like atoms*, Phys. Scr. **70** (2004), 276–279.
- [49] M. L. Sanchez, B. Moreno, and A. López Piñeiro, *Matrix-element calculations for hydrogenlike atoms*, Phys. Rev. A **46** (1992) #11, 6908–6913.
- [50] V. M. Shabaev, *Recurrence formulas and some exact relations for radial integrals with Dirac and Schrödinger wave functions*, Vestnik Leningradskogo Universiteta, Seria 4: Fizika, Khimiya (1984) number 1, 15–19 [in Russian].
- [51] V. M. Shabaev, *Generalizations of the virial relations for the Dirac equation in a central field and their applications to the Coulomb field*, J. Phys. B: At. Mol. Opt. Phys. **24** (1991), 4479–4488.
- [52] V. M. Shabaev, *Relativistic recoil corrections to the atomic energy levels*, in: *The Hydrogen Atom: Precision Physics of Simple Atomic Systems*, Lecture Notes in Physics, Vol. 570, Springer-Verlag, Berlin, Heidelberg, 2001, pp. 714–726.
- [53] V. M. Shabaev, *Two-time Green's function method in quantum electrodynamics of high-Z few-electron atoms*, Phys. Rep. **356** (2002), 119–228.
- [54] V. M. Shabaev, *Virial relations for the Dirac equation and their applications to calculations of hydrogen-like atoms*, in: *Precision Physics of Simple Atomic Systems*, Lecture Notes in Physics, Vol. 627, Springer-Verlag, Berlin, Heidelberg, 2003, pp. 97–113.
- [55] V. M. Shabaev, *Quantum electrodynamics of heavy ions and atoms: current status and prospects*, Phys.-Uspekhi **178** (2008) #11, 1220–1225 [in Russian]; Phys.-Uspekhi **51** (2008) #11, 1175–1180.
- [56] J. Shertzer, *Evaluation of matrix elements $\langle n, l | r^\beta | n, l' \rangle$ for arbitrary β* , Phys. Rev. A **44** (1991) #5, 2832–2835.
- [57] S. K. Suslov, *Expectation values in relativistic Coulomb problems*, J. Phys. B: Atom. Mol. Opt. Phys. **42** (2009), 185003 (8pp); published on line 9 September 2009, URL: <http://www.iop.org/EJ/abstract/0953-4075/42/18/185003/>.
- [58] S. K. Suslov and B. Trey, *The Hahn polynomials in the nonrelativistic and relativistic Coulomb problems*, J. Math. Phys. **49** (2008) #1, 012104 (51pp); published on line 22 January 2008, URL: <http://link.aip.org/link/?JMP/49/012104>.
- [59] R. A. Swainson and G. W. F. Drake, *An alternative proof of some relations between hydrogenic matrix elements*, J. Phys. B: At. Mol. Opt. Phys. **23** (1990) #7, 1079–1084.
- [60] P. L. Tchebychef, *Sur l'interpolation par la méthode des moindres carrés*, Mémoires de l'Académie Impériale des sciences de St.-Petersbourg, VII^e serie, **1** (1859) #15, 1–24; reprinted in: *Œuvres*, volume 1, Chelsea Publishing Company, New York, 1962, pp. 473–498.
- [61] P. L. Tchebychef, *Sur l'interpolation*, (1864); reprinted in: *Œuvres*, volume 1, Chelsea Publishing Company, New York, 1962, pp. 542–560.
- [62] P. L. Tchebychef, *Sur l'interpolation des valeurs équidistantes*, (1875); reprinted in: *Œuvres*, volume 2, Chelsea Publishing Company, New York, 1962, pp. 219–242.
- [63] J. H. Van Vleck, *A new method of calculating the mean value of $1/r^s$ for Keplerian systems*, Proc. Roy. Soc. London A **143** (1934), 679–681.
- [64] E. R. Vrscaj and H. Hamidian, *Rayleigh–Schrödinger perturbation theory at large order for radial relativistic Hamiltonians using hypervirial Hellmann–Feynman theorems*, Phys. Lett. A **130** (1988) #3, 141–146.
- [65] I. Waller, *Der Starkeffekt zweiter Ordnung bei Wasserstoff und die Rydbergkorrektur der Spektren von He und Li⁺*, Zeitschrift für Physik **38** (1926), 635–646.

SCHOOL OF MATHEMATICAL AND STATISTICAL SCIENCES AND MATHEMATICAL, COMPUTATIONAL AND MODELING SCIENCES CENTER, ARIZONA STATE UNIVERSITY, TEMPE, AZ 85287-1804, U.S.A.

E-mail address: `sks@asu.edu`

URL: `http://hahn.la.asu.edu/~suslov/index.html`